

**Patient  
Reminder Card**

**Remsima<sup>®</sup>**  
Infliximab

**Show this card to  
any doctor involved  
in your treatment.**

Name patient:

.....  
.....

Name doctor:

.....  
.....

Telephone number doctor:

.....  
.....

Date of Remsima therapy  
initiation:

.....  
.....

Current administrations:

.....  
.....

Please make sure you also  
have a list of all other  
medicines that you are using  
with you at any visit to a  
healthcare professional.

List of other medicines:

.....  
.....

List of allergies:

.....  
.....

Ask your doctor to record  
the type and date of last  
screening(s) for tuberculosis  
(TB) below:

Test:

Date:

Result:

It is important that you and  
your doctor record the  
brand name and batch  
number of your medicine.

Brand name:

Batch number:

.....



LM141007-PA001

**This Patient Reminder Card contains important safety information that you need to be aware of before and during treatment with Remsima.**

When starting a new card, please keep this card as a reference for 4 months after your last dose of Remsima.

Please read the Remsima 'Package Leaflet' carefully before you start using this medicine.

Keep this card with you for 4 months after your last dose of Remsima, or in case of pregnancy, for at least 6 months after the birth of your baby.

Side effects may occur a long time after your last dose.

**| Infections |**  
**Before treatment with Remsima**

- Tell your doctor if you have an infection even if it is a very minor one.
- It is very important that you tell your doctor if you have ever had tuberculosis (TB), or if you have been in close contact with someone who has had TB. Your doctor will test you to see if you have TB. Ask your doctor to record the type and date of your last screening(s) for TB on the card.

- Tell your doctor if you have hepatitis B or if you know or suspect you are a carrier of the hepatitis B virus.

**During treatment with Remsima**

- Tell your doctor straight away if you have signs of an infection. Signs include a fever, feeling tired, (persistent) cough, shortness of breath, weight loss, night sweats, diarrhoea, wounds, dental problems, burning when urinating or 'flu-like' signs.

**| Pregnancy and Vaccinations |**

- In case you have received Remsima while you were pregnant, it is important that you inform your baby's doctor about it before your baby receives any vaccine. Your baby should not receive a 'live vaccine', such as BCG (used to prevent tuberculosis) within 6 months after birth.